



Marlow, R. D., Finn, A., & Henderson, J. (2019). Assessing the association between bronchiolitis in infancy and recurrent wheeze: A whole English birth cohort case-control study. *Thorax*, 74(5), 503-505. <https://doi.org/10.1136/thoraxjnl-2018-212203>

Peer reviewed version

License (if available):
CC BY-NC

Link to published version (if available):
[10.1136/thoraxjnl-2018-212203](https://doi.org/10.1136/thoraxjnl-2018-212203)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via BMJ Publishing at <https://thorax.bmj.com/content/74/5/503> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Assessing the Association between Bronchiolitis in Infancy and Recurrent Wheeze – A Whole English Birth Cohort Case Control Study.

Robin Marlow¹, Adam Finn¹, John Henderson¹

1) Population Health Sciences, University of Bristol, UK

Address for correspondence:

Dr Robin Marlow, Level 6, UH Bristol Education and Research Centre,

Upper Maudlin Street Bristol BS2 8AE

Email: robin.marlow@bristol.ac.uk

Tel: 0117 3420172

Fax: [0117 342 0209](tel:01173420209)

Keywords

Asthma Epidemiology

Clinical Epidemiology

Respiratory Infection

Word count - 1327

Abstract

The precise association between bronchiolitis and predisposition to childhood wheeze is unclear. We assessed bronchiolitis aetiology and later wheeze phenotypes in the entire 2007 English birth cohort. All infants admitted to hospital in England during their first year of life with bronchiolitis or urinary tract infection (UTI) were followed using Hospital Episode Statistics, to determine risk and characteristics of wheeze admission over the subsequent 8 years. In our cohort of 21,272 children compared to UTI, the risk of wheeze admission was higher with previous bronchiolitis (RR2.4), even higher in those with non-RSV bronchiolitis (RR3.1) and persisted into late-onset wheeze (RR1.7).

1 Introduction

2 Bronchiolitis is the commonest reason for hospital admission in infancy. In the UK, currently four
3 percent of the children born per year are admitted to hospital with bronchiolitis (40.4/1000).[1]
4 Wheeze is one of the commonest causes for admission in later childhood – 3.1 admissions / 1000
5 children in 2016.[2] These two respiratory diagnoses combined represent a significant burden for
6 both children, their families and the health service.[3] The link between respiratory syncytial virus
7 (RSV) bronchiolitis and early (post bronchiolitis) wheeze is recognised, however the association with
8 late phenotypes of wheeze and other viruses is less clear.[4] With potential vaccines and
9 therapeutic agents in development against RSV; accurately quantifying any potentially causal link
10 could be of use for their health economic assessment. In this study we sought to characterise the
11 association between different viral causes of bronchiolitis and phenotypes of childhood wheeze.

12 Method

13 Hospital Episode Statistics discharge codes were used to create two cohorts; all infants (<12m of
14 age) admitted to hospital in England during 2007 diagnosed with bronchiolitis or as a comparator
15 with a urinary tract infection (UTI - chosen as the commonest non-respiratory cause for infant
16 admission to hospital). [2] Our case definition of bronchiolitis was children <12 months of age
17 admitted to hospital with a clinician diagnosis of bronchiolitis (table 1). The all-cause bronchiolitis
18 (acBr) cohort was subdivided by coded causative organism (RSV, non RSV (nRSV) or un-specified
19 bronchiolitis (usBr)). Patients with both diagnoses during the first year of life were excluded as were
20 all those who had been admitted to neonatal intensive care at birth (due to an increased risk of
21 chronic lung disease). From their first birthday we compared the number and pattern of admissions
22 with wheeze between the two cohorts over the subsequent eight years. Timings of recurrent
23 wheeze episodes were used to categorise patients into previously described wheeze phenotypes.[5]
24 To test for confounding between our cohorts we compared the rates of head injury (a common non-
25 infective reason for paediatric admission) as a negative control, with the hypothesis that if unbiased
26 that this should not differ between cohorts.[6] R v3.4.1 was used for data cleaning and to perform
27 log-binomial regression adjusted for gender to calculate relative risks.

28 Results

29 From the 2007 birth cohort of 642,000 children, we identified 17,946 cases of bronchiolitis and 3,326
30 of UTI (Table 1) with risks of later admission with any wheeze of 19% & 7 % respectively; RR of
31 admission 2.4 (95%CI 2.2-2.8). As there was a significant difference in gender between the cohorts,
32 60% male in bronchiolitis vs 48% male in UTI we adjusted for gender in our models. Our negative
33 control suggested no bias with no difference found in the risk of head injury admission (RR 0.96,
34 95%CI 0.7-1.3) between the cohorts.

Against all cause wheeze admissions, nRSV bronchiolitis had a significantly stronger association RR 3.1(2.4-3.9) than for RSV RR 2.3 (2.0-2.6) $p=0.01$ but not the undifferentiated usBr RR 2.5 (2.2-2.8) $p=0.06$. Using length of bronchiolitis admission as a proxy for disease severity there was an apparent dose effect, with risk of later wheeze increased with admissions <24hrs and steadily increasing for longer admissions (table 2). All cause bronchiolitis increased the risk of all phenotypes of wheeze, particularly Transient, Persistent Early and Persistent. (Table 3)

Discussion.

We have shown that significantly more cases of all phenotypes of wheeze occur in children that have had bronchiolitis in infancy compared to those who had a UTI, with the largest effects seen in long admissions with non-RSV bronchiolitis. This effect is most marked for the early wheeze phenotypes but persists for even late onset wheeze.

Whilst the link between RSV bronchiolitis and later respiratory pathology has been consistently demonstrated, the strength and details of the association vary between populations and has not been previously described in a UK cohort.[7] Studies have also linked rhinovirus, human metapneumovirus and parainfluenzae as associated with later wheeze, with some suggesting a stronger effect for non-RSV bronchiolitis.[8] The strongest form of evidence for assessing the association between bronchiolitis and wheeze is from randomised controlled trials (RCT) of monoclonal antibodies against RSV. The Dutch MAKI RCT of Paluvizumab in premature infants found at 6 years there was a significant reduction in parentally reported wheeze (11.6% vs 19.9%) but not in physician diagnosed (10.3% vs 9.9%), concluding that there was no clinically significant effect.[9] The phase 3 RCT of Motavizumab in 2170 term Navajo infants found an 87% relative reduction in RSV related admissions but no effect on rates of medically attended wheezing (14.9% vs 14.0%) during 1–3 years follow-up.[10] In our cohorts rates of bronchiolitis associated late onset wheeze were low (5% vs 3%); raising the possibility that with <1000 patients per arm and only a subset of these developing bronchiolitis, these trials were not adequately powered to detect this difference. Although a small percentage change, we would argue that on a population scale a vaccine that prevented any cases of a chronic health condition such as asthma as a *secondary* gain would be significant.

The strengths of this paper are that it follows a whole English population birth cohort of 642,000 children over an 8 year period. Using a hard outcome of admission rather than parentally reported or clinician diagnosed wheeze it characterises the burden of admissions due to different phenotypes of wheeze. Our primary analysis using all diagnostic fields found similar frequencies of wheezing during admission as in parentally reported community surveys,[11] restricting analysis to the

primary diagnostic code found reduced incidence of wheeze (7.8% post bronchiolitis vs 3.1% post UTI) but with a similar risk ratio 2.4 (2.0-3.0) between the groups. Its limitations are that it is reliant on quality of diagnostic coding, rates of routine virological testing and could not assess the effects on mild non-admitted wheeze presentations. As a retrospective analysis of routine health data we were not able to distinguish between different syndromic phenotypes of bronchiolitis as our dataset limited us to identifying patients by diagnostic clinical coding. Rates of routine virological testing carried out in the UK in 2007 meant we could only meaningfully compare RSV to non-RSV subgroups. Our cohort of hospitalised children represent a severe subgroup and thus our findings are likely to be an overestimate of true population effect. Our analysis was controlled for gender, in a sensitivity analysis we also included the Index of Multiple Deprivations but this did not alter the relative risk estimates. There may be other biases that we were not able to identify. Fundamentally as an observational study design we cannot hope to prove causality and must be cautious that our findings are not a product of the phenotype categorisation.[12]

Whilst the narrative that bronchiolitis leads to an inflammatory state, the start of the allergic march and predisposition to later wheeze is appealing; that all causes of bronchiolitis are associated with all phenotypes of wheeze suggests this interpretation is actually the cart leading the horse. Perhaps bronchiolitis is simply a marker of children who are likely to go on to wheeze in later life and represents the sentinel manifestation of those genetically or environmentally pre-disposed to wheeze. That the association for early wheeze is stronger implies that this is more complicated than the revealing of an innate state. It may be that different clinical phenotypes of bronchiolitis[13] would be helpful in this prognostication but we were not able to assess this using this dataset.

Fundamentally these data highlight our inability to prospectively clinically distinguish between wheeze phenotypes. It is reassuring that 80% of children with even severe bronchiolitis do not go on to wheeze. Academic interest aside, the reason for this study is to be able to answer parents' questions of "will this happen again?" Although these data cannot elucidate the cause, they are helpful to describe likely respiratory trajectories for children. We now plan to expand this research by also examining wheeze attendances in the emergency department and primary care to see if the association still holds for the less severe cases of bronchiolitis.

HES data statement – The School of Social and Community Medicine (SSCM), University of Bristol, has a Data Sharing Agreement (DSA; NIC-1785-X7K1V) with the HSCIC for HES Admitted Patient Care (inpatient / day case) data for the financial years 2005/6 to 2014/15. The purchase of these data was funded by NIHR CLAHRC West. Copyright © 2018, re-used with the permission of The Health & Social Care Information Centre. All rights reserved. Our data access agreement with HSIC does not allow

101 distribution of source data but extraction and analysis scripts are available on request. This work
102 uses data provided by patients and collected by the NHS as part of their care and support and would
103 not have been possible without access to this data. The NIHR recognises and values the role of
104 patient data, securely accessed and stored, both in underpinning and leading to improvements in
105 research and care.

106

107 **Tables**

Groups		Number	ICD-10 codes
Bronchiolitis (60% male)	RSV	4399	J12.1, J21.0,
	Non-RSV	243	J12.0, J12.2, J12.3, J12.8, J21.1, J21.8,
	Unspecified	13304	J12.9, J21.9
UTI (control) (52% male)		3326	N39.0
Outcomes	Wheezing	4472	J45.*, J46*, R06.2
	Head Injury	325	S09*
Excluded:	Neonatal admissions	4654	n/a
	Both diagnoses	191	n/a

108 Table 1 – Size of cohorts and defining ICD10 codes.

109

Length of admission (days)	0	1->3	4->6	7-13	>14	Combined
All Bronchiolitis RR (95% CI)	2.16 (1.9-2.5)	2.37 (2.1-2.7)	2.57 (2.3-2.9)	3.37 (2.9-3.9)	3.24 (2.4-4.3)	2.4 (2.2-2.8)
RSV RR (95% CI)	1.6 (1.2-2.2)	2.0 (1.7-2.4)	2.3 (1.7-2.4)	3.1 (2.6-3.7)	2.9 (1.8-4.2)	2.3 (2.0-2.6)
nRSV RR (95% CI)	2.5 (0.8-5.2)	1.5 (0.7-2.6)	3.5 (2.3-4.9)	5 (3.3-7.0)	3.6 (1.4-6.6)	3.1 (2.4-4.0)
usBR RR (95% CI)	2.2 (1.9-2.5)	2.5 (2.2-2.8)	2.7 (2.4-3.1)	3.5 (2.9-4.3)	3.7 (2.3-5.5)	2.5 (2.2-2.8)

110 Table 2 –risk ratio of later admission with wheeze by aetiology and length of bronchiolitis admission.

111

Wheeze	All Wheeze	Phenotype					
		NEVER	TREY	PEEY	INTER	LATE	PRST
Bronchiolitis	18.7%	80.1%	4.4%	0.8%	7.6%	5.0%	1.0%
UTI	7.2%	90.4%	0.9%	0.2%	3.1%	3.1%	0.2%
RR	2.43	n/a	5.4	4.8	2.6	1.7	5.2
95% CI	2.2-2.8	n/a	3.8-8.0	2.3- 12.1	2.1-3.1	1.4-2.1	2.6- 12.2

112 Table 3 - Distribution of wheeze phenotypes for different cohorts and resulting risk ratios.

113 (Phenotypes: NEVER - no wheeze;

114 TREY - Transient early, wheeze before 18m until 42m;

115 PEEY – Prolonged Early, wheeze before 18 until 69m;

116 INTER – Intermediate, no wheeze until after 42m;

117 LATE – Late onset, no wheeze until after 69m;

118 PRST – Persistent, wheeze before 18m until >69m
119) [5]

120

121 **Contribution**

122 RM study design, analysis and authorship of first draft. AF, JH critically appraised and
123 developed the manuscript.

124

125 **Funding**

126 RM is funded by the NIHR as an Academic Clinical Lecturer and is a member of the NIHR
127 Health Protection Research Unit in Evaluation of Interventions at University of Bristol.

128 **Competing Interests**

129 No competing interests

130 **Exclusive licence**

131 The Corresponding Author has the right to grant on behalf of all authors and does grant on
132 behalf of all authors, an exclusive licence (or non-exclusive for government employees) on a
133 worldwide basis to the BMJ Publishing Group Ltd and its Licensees to permit this article (if
134 accepted) to be published in Thorax and any other BMJ PGL products to exploit all subsidiary
135 rights, as set out in our licence [http://journals.bmj.com/site/authors/editorial-](http://journals.bmj.com/site/authors/editorial-policies.xhtml#copyright)
136 [policies.xhtml#copyright](http://journals.bmj.com/site/authors/editorial-policies.xhtml#copyright) and the Corresponding Author accepts and understands that any
137 supply made under these terms is made by BMJ PGL to the Corresponding Author. Where
138 the Corresponding Author wishes to make the article available on an open access basis (and
139 intends to pay the relevant open access fee), the terms of such open access shall be
140 governed by a Creative Commons licence – details of these licences and which Creative
141 Commons licence will apply to this article are set out in our licence referred to above.

142 I confirm this manuscript has not been accepted for publication elsewhere, is not being
143 considered for publication elsewhere and does not duplicate material already published. I
144 confirm all authors consent to publication of this manuscript.

References

- 1 Green CA, Yeates D, Goldacre A, *et al.* Admission to hospital for bronchiolitis in England: trends over five decades, geographical variation and association with perinatal characteristics and subsequent asthma. *Arch Dis Child* 2016;**101**:140–6. doi:10.1136/archdischild-2015-308723
- 2 Hospital Admitted Patient Care Activity, 2016-17. NHS Digit. <https://digital.nhs.uk/data-and-information/publications/statistical/hospital-admitted-patient-care-activity/2016-17> (accessed 23 Aug 2018).
- 3 Stevens CA, Turner D, Kuehni CE, *et al.* The economic impact of preschool asthma and wheeze. *Eur Respir J* 2003;**21**:1000–6. doi:10/cdvfdw
- 4 Régnier SA, Huels J. Association between respiratory syncytial virus hospitalizations in infants and respiratory sequelae: systematic review and meta-analysis. *Pediatr Infect Dis J* 2013;**32**:820–6. doi:10.1097/INF.0b013e31829061e8
- 5 Henderson J, Granell R, Heron J, *et al.* Associations of wheezing phenotypes in the first 6 years of life with atopy, lung function and airway responsiveness in mid-childhood. *Thorax* 2008;**63**:974–80. doi:10.1136/thx.2007.093187
- 6 Lipsitch M, Tchetgen Tchetgen E, Cohen T. Negative controls: a tool for detecting confounding and bias in observational studies. *Epidemiol Camb Mass* 2010;**21**:383–8. doi:10/drzms4
- 7 Fauroux B, Simões EAF, Checchia PA, *et al.* The Burden and Long-term Respiratory Morbidity Associated with Respiratory Syncytial Virus Infection in Early Childhood. *Infect Dis Ther* 2017;**6**:173–97. doi:10/gc4ng9
- 8 Koponen P, Helminen M, Paasilta M, *et al.* Preschool asthma after bronchiolitis in infancy. *Eur Respir J* 2012;**39**:76–80. doi:10/c74d4n
- 9 Mochizuki H, Kusuda S, Okada K, *et al.* Palivizumab Prophylaxis in Preterm Infants and Subsequent Recurrent Wheezing. Six-Year Follow-up Study. *Am J Respir Crit Care Med* 2017;**196**:29–38. doi:10/gbm6t
- 10 O'Brien KL, Chandran A, Weatherholtz R, *et al.* Efficacy of motavizumab for the prevention of respiratory syncytial virus disease in healthy Native American infants: a phase 3 randomised double-blind placebo-controlled trial. *Lancet Infect Dis* 2015;**15**:1398–408. doi:10.1016/S1473-3099(15)00247-9
- 11 Kuehni CE, Strippoli M-PF, Low N, *et al.* Wheeze and asthma prevalence and related health-service use in white and south Asian pre-schoolchildren in the United Kingdom. *Clin Exp Allergy J Br Soc Allergy Clin Immunol* 2007;**37**:1738–46. doi:10/fvx6d2
- 12 Richiardi L, Barone-Adesi F, Pearce N. Cancer subtypes in aetiological research. *Eur J Epidemiol* 2017;**32**:353–61. doi:10.1007/s10654-017-0253-z
- 13 Dumas O, Mansbach JM, Jartti T, *et al.* A clustering approach to identify severe bronchiolitis profiles in children. *Thorax* 2016;**71**:712–8. doi:10.1136/thoraxjnl-2016-208535